

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 428 941 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(43) Date of publication of patent specification: 03.05.95 (51) Int. Cl.⁶: C07D 251/04, A01N 43/64

(21) Application number: 90121383.5

(22) Date of filing: 08.11.90

(54) Hexahydrotriazine compounds and insecticides.

(30) Priority: 10.11.89 JP 292675/89
02.02.90 JP 24199/90

(43) Date of publication of application:
29.05.91 Bulletin 91/22

(48) Publication of the grant of the patent:
03.05.95 Bulletin 95/18

(84) Designated Contracting States:
CH DE FR GB IT LI

(56) References cited:
EP-A- 0 308 961
DD-A- 155 030

CHEMICAL ABSTRACTS, vol. 111, no. 18, October 18, 1989, Columbus, Ohio, USA YU, YONGZHONG et al. "Synthesis of polynitro compounds from nitroguanidine" page 195, column 2, abstract no. 157 006q & Propellants, Explos., Pyrotech. 1989, 14(4), 150-2

CHEMICAL ABSTRACTS, vol. 97, no. 3, July 19, 1982, Columbus, Ohio, USA YAO, GUOWEI et al. : "Synthesis and reactions of N-(2,2-dinitroethyl)-N'-nitroguanidine", page 709, column 1, abstract-no. 23746q & Kogyo Kayaku 1982, 43(1), 2-8

(73) Proprietor: AGRO-KANESHO CO., LTD.
No. 4-1, Marunouchi 2-chome
Chiyoda-ku
Tokyo (JP)

(72) Inventor: Wu, Frank
1-307-1-207, Noshio
Kiyose-shi,
Tokyo (JP)
Inventor: Kariya, Akinori
3-13-18, Hagiyama-cho
Higashi-Murayama-shi,
Tokyo (JP)
Inventor: Katsuyama, Noriyoshi
2-5-5, Hachiman,
Kawajima-machi
Hiki-gun,
Saitama-ken (JP)
Inventor: Tsuji, Atsushi
1-36-3, Kitazawa,
Setagaya-ku
Tokyo (JP)
Inventor: Takasuka, Kiyoshi
Green-Copo 207,
1-632-45, Higashi-Sayamagaoka
Tokorozawa-shi,
Saitama-ken (JP)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 428 941 B1


CHEMICAL ABSTRACTS, vol. 109, no. 6, August 8, 1988, Columbus, Ohio, USA PER-
NYESZI, JOSEF et al. "Revision of manufacturing process of
1,3,5-tris(2-chlorophenyl)hexahydro-1,3,5-
triazine for reasons of economy and environmental protection" page 104, column 1,
abstract -no. 39 691v & Nehezvegylp. Kut.
Intez. Kozl. 1986, 17, 155-70

Inventor: Segami, Shigenori
Haimu H.T. 2-203,
137, Yamaguchi
Tokorozawa-shi,
Saltama-ken (JP)
Inventor: Nanjo, Katsumi
1563-7, Shimoyasumatsu
Tokorozawa-shi,
Saltama-ken (JP)
Inventor: Sato, Junko
1-17-30-201, Nishi
Kunitachi-shi,
Tokyo (JP)


Ⓓ Representative: VOSSIUS & PARTNER
Postfach 86 07 67
D-81634 München (DE)

Description

The present invention relates to hexahydrotriazine compounds, a process for preparing the same and insecticides containing the same as active ingredients. The hexahydrotriazine compounds provided by this invention are represented by the following general formula (I):



wherein R stands for lower alkyl or lower alkenyl; R₁ stands for lower alkyl, lower alkenyl or lower alkynyl; and R₂ stands for a group indicated by




A variety of chemicals have been developed and used as insecticides for controlling harmful insects. Representative examples of such chemicals are organic phosphorous compounds, carbamate compounds and synthetic pyrethroids. However, by the repeated use of these insecticides, many harmful insects have acquired resistance to these chemicals, thus making it difficult to control such insects. On the other hand, although some conventional insecticides have high insecticidal activity, they might cause environmental pollution problems either due to their high toxicity to warm-blooded animals or fishes and Crustacea or due to their remaining in the environment in large amounts for a long time after use, thus putting the ecological system out of order. Accordingly, there is a demand for the development of novel insecticides which more effectively control harmful insects, which have acquired resistance to conventional chemicals, and yet are low in toxicity not only to warm-blooded animals but also fishes and Crustacea. There is also a demand for novel insecticides which remain in plants and soils in decreased amounts after their use, and do not produce phytotoxicity in various plants.

An object of this invention is to provide novel compounds which (1) more effectively control harmful insects that have acquired resistance to conventional chemicals, (2) are low in toxicity not only to warm-blooded animals but also fishes and Crustacea and (3) remain in decreased amounts in plants and soils after their use. Lastly, such novel compounds do not produce phytotoxicity in various plants.


Another object of this invention is to provide a process for preparing such compounds.

A further object of this invention is to provide improved insecticides which (1) more effectively control harmful insects having resistance to conventional insecticides, (2) are low in toxicity not only to warm-blooded animals but also fishes and Crustacea, and (3) remain in decreased amounts in plants and soils after their use. Lastly, such improved insecticides do not produce phytotoxicity in various plants.

These objects could be achieved by the provision of the new hexahydrotriazine compounds represented by the following general formula (I):



wherein R stands for lower alkyl or lower alkenyl; R₁ stands for lower alkyl, lower alkenyl or lower alkynyl; and R₂ stands for a group indicated by




which can be used as extremely powerful insecticides which exhibit rapid effects.


The term "lower" in connection with alkyl, alkenyl and alkynyl generally comprises straight and branched chain residues having 1 to 8, preferably 1 to 6 most preferably 1 to 4 carbon atoms. Specific examples are the methyl, ethyl, propyl, isopropyl and allyl groups.

The compounds of this invention represented by the general formula (I) may be prepared by the processes described below.

Preparation Process (a)



wherein R stands for lower alkyl or lower alkenyl; R₁ stands for lower alkyl, lower alkenyl, lower-alkynyl; R₂ stands for a group indicated by




and X stands for halogen.

In the preparation process (a), as indicated by the reaction formula set forth above, the compounds of this invention may be readily prepared by reacting, in a solvent, a compound represented by the general formula (II) with a compound represented by the general formula (III), such as bis(chloromethyl)-methylamine, bis(chloromethyl)ethylamine and bis(chloromethyl)propylamine, in the presence of a basic material.

Examples of the suitable solvents which may be used in this reaction are ethers such as diethyl ether and tetrahydrofuran; acetonitrile; aromatic hydrocarbons; chlorinated hydrocarbons; DMF; and DMSO. These solvents may be used alone or in combination. Compounds which may be used as the basic materials are, for example, inorganic bases such as sodium hydroxide and potassium carbonate, and organic bases such as pyridine and triethylamine. These bases may be used in excess and particularly, 2.1 to 4 mols per 1 mol of the compound (II) are preferable.

The reaction temperature may be freely set within a range of from -20 °C to 100 °C, preferably from -5 °C to 10 °C.


Preparation Process (b)



wherein R, R₁ and R₂ have the same meanings defined above.


In the preparation process (b) as indicated by the reaction formula set forth above, the compounds of this invention may be readily prepared by reacting a compound represented by the general formula (II) with a primary amine, represented by the general formula R-NH₂ (IV) wherein R has the same meaning defined above, such as methyl amine, ethyl amine and allyl amine and a 37% formalin solution or paraformaldehyde, in a solvent.

Examples of the suitable solvents which may be used in this reaction are: alcohols such as methanol and ethanol; ethers such as dioxane and tetrahydrofuran; and water. These solvents may be used alone or in combination. The reaction temperature may be freely set within a range from a room temperature to a boiling temperature of a solvent used, preferably from 50 °C to 100 °C. Preparation Process (c)



wherein R, R₁ and R₂ have the same meanings defined above and Y stands for halogen or a group indicated by -OSO₂CH₃ or

5



10 In the preparation process (c), as indicated by the reaction formula set forth above, the compounds of this invention may be readily prepared by reacting a compound represented by the general formula (V) with a compound represented by the general formula (VI) in the presence of a basic material in an inert solvent.

Examples of the suitable solvents which may be used in this reaction are: DMF; DMSO; acetonitrile; and ethers, such as tetrahydrofuran and dioxane. These solvents may be used alone or in combination.
 15 Compounds which may be used as the basic materials are for example, sodium hydride, potassium carbonate, sodium hydroxide etc. 1.1 to 2.5 mols of these bases per 1 mol of the compound (V) are preferably used. The reaction temperature may be freely set within a range from a room temperature to 150 °C, preferably from 40 °C to 100 °C.

Hexahydrotriazine compounds of this invention are novel compounds. Typical compounds of this
 20 invention are set forth in the following Table 1.

25

30

35


40

45

50

55

Table 1








Compound No.	R ₁	R ₂	R	Melting Point
1 (Reference)		H	C ₆ H ₅	150~154 °C
2 (Reference)	"	"	C ₆ H ₅	124~125 °C
3 (Reference)	"	"	i C ₆ H ₅	121~122 °C
5	"	C ₆ H ₅	C ₆ H ₅	116~117 °C
6		"	"	125~126 °C
7		C ₆ H ₅	"	163~164 °C
8	"	n C ₆ H ₅	"	102~104 °C

Table 1 (continued)

Compound No.	R ₁	R ₂	R	Melting Point
9		n C ₂ H ₅	CH ₃	139~143 °C
10		-CH ₂ CH=CH ₂	"	112~114 °C
11	"	-CH ₂ C≡CH	"	124~125 °C

An effective amount of any of the compounds of this invention may be applied directly as an insecticide or may be formulated by conventional technology in the form of an emulsion, a wettable powder, a dust, a granule or in a flowable form, and then applied as a formulated product. A liquid or solid carrier may be used for preparing a formulated product containing one or more of the compounds of this invention. Liquid carriers which may be used for this purpose include, for example, organic solvents, and conveniently used are xylene, chlorobenzene, methylnaphthalene, cyclohexanone, isophorone, alcohols, dimethylformamide and N-methylpyrrolidone. Examples of solid carriers include kaoline, talc, bentonite, diatomaceous earth and clay, and synthetic compounds such as alumina, zeolite and silicates may also be used. In preparation of the formulated products, various adjuvants, such as emulsifiers, dispersants, spreaders, wetting agents and penetrating agents, may be used for providing the products with the desired properties of emulsification, dispersion, suspension and penetration.

The compounds of this invention, which are represented by the general formula (I) set forth above, have strong insecticidal activity against various harmful insects, including Hemiptera, Lepidoptera, Coleoptera, Diptera, Orthoptera and Isoptera, and yet are low in toxicity to human beings and animals and do not produce phytotoxicity in various plants, and thus can be practically used as superior insecticides.

Examples of harmful insects which may be effectively controlled by the compounds of this invention are as follows:

Hemiptera such as *Nilaparvata lugens*, *Laodelphax striatellus*, *Nephotettix cincticeps*, *Pseudococcus comstocki*, *Unaspis yanonensis*, *Myzus persicae*, *Aphis gossypii*, *Lipaphis erysimi*, *Stephanitis nashi*, *Scotinophara lurida* and *Trialeurodes vaporariorum*; Lepidoptera such as *Pieris rapae*, *Spodoptera litura*, *Mamestra brassicae*, *Chilo suppressalis*, *Plutera xylostella*, *Adoxophyes orana*, *Agrotis segetum*, *Cnaphalocrocis medinalis* and *Ostrinia furnacalis*; Coleoptera such as *Henosepilachna vigintioctopunctata*, *Aulacophora femoralis*, *Phyllotreta striolata*, *Lissorhoptrus oryzophilus*, *Sitophilus zeamais* and *Anomala rufocuprea*; Diptera such as *Musca domestica*, *Hylemia platura*, *Culex pipiens*; Orthoptera such as *Gryllotalpa africana*, *Blatella germanica* and *Locusta migratoria*; and Isoptera such as *Coptotermes formosanus* and *Reticulitermes speratus*.

The present invention will now further be explained by way of the following non-limiting examples.

Examples

The processes for the preparation of reference compounds and of the compounds of this invention are described in detail in the following Synthesis Examples.

Synthesis Example 1 (reference)

Synthesis of 1-(2-chloro-5-pyridylmethyl)-5-methyl-2-nitroimino-hexahydro-1,3,5-triazine

0.6 g of 1-(2-chloro-5-pyridylmethyl)-3-nitroguanidine and 0.4 g of bis(chloromethyl)methylamine were suspended in dried THF. To the suspension, a solution of 0.52 g of triethylamine in THF was added

dropwise with cooling. The reaction mixture was stirred for 1 hour, and then poured into ice water. The mixture was extracted with dichloromethane. After the extract was dried over anhydrous magnesium sulfate, dichloromethane was distilled off. The residue was recrystallized from methanol to obtain 0.78 g of the title compound. This compound is No. 1 in Table 1. m.p.: 150 - 154 °C

5

Synthesis Example 2 (reference)

Synthesis of 1-(2-chloro-5-pyridylmethyl)-5-ethyl-2-nitroimino-hexahydro-1,3,5-triazine

10 To a stirring mixture of 1 g of 1-(2-chloro-5-pyridylmethyl)-3-nitroguanidine, 0.28 g of 70% ethylamine in water and 20 ml of ethanol, 0.71 g of 37% formalin solution was added dropwise with heating. The reaction mixture was heated under reflux for two hours. After being allowed to cool to room temperature, the crystals formed, which were filtrated off and recrystallized from methanol to obtain 1.15 g of the title compound. This compound is No. 2 in Table 1. m.p.: 124 - 125 °C

15

Synthesis Example 3

Synthesis of 1-(2-chloro-5-pyridylmethyl)-3,5-dimethyl-2-nitroimino-hexahydro-1,3,5-triazine

20 1 g of 1,5-dimethyl-2-nitroimino-hexahydro-1,3,5-triazine was dissolved in 20 ml of dried DMF. To the solution, 0.27 g of 60% sodium hydride was added with cooling. The mixture was stirred for 1 hour at room temperature until evolution of hydrogen was ceased and then the mixture was heated with stirring further for 1 hour at 50 °C. To the mixture, a solution of 0.9 g of 2-chloro-5-pyridylmethylchloride in 8 ml of dried DMF was added dropwise at 40 - 50 °C. After this addition, the reaction mixture was heated with stirring it for two
25 hours at 70 - 80 °C. The mixture was poured into ice-water and extracted with dichloromethane. The extract was dried over anhydrous magnesium sulfate, and dichloromethane was distilled off. The residue was purified by a column chromatography to obtain 1.3 g of the title compound. This compound is No.5 in Table 1. m.p.: 116 - 117 °C

Synthesis Example 4

Synthesis of 1-(2-chloro-5-thiazolylmethyl)-3,5-dimethyl 2-nitroimino-hexahydro-1,3,5-triazine

35 1.7 g of 1,5-dimethyl-2-nitroimino-hexahydro-1,3,5-triazine was dissolved in 20 ml of dried DMF. To the solution, 0.28 g of 60% sodium hydride was added portionwise with cooling. The mixture was stirred for 1 hour at room temperature until evolution of hydrogen was ceased and then the mixture was heated with stirring further for 1 hour at 50 °C. To the mixture, a solution of 1.7 g of 2-chloro-5-thiazolylmethylchloride in 8 ml of dried DMF was added dropwise at 40 - 50 °C. After this addition, the reaction mixture was heated with stirring for two hours at 70 - 80 °C. The mixture was poured into ice-water and extracted with
40 dichloromethane. The extract was dried over anhydrous magnesium sulfate, and dichloromethane was distilled off. The residue was purified by a column chromatography to obtain 0.82 g of the title compound. This compound is No.6 in Table 1. m.p.: 125 - 126 °C

Some specific Formulation Examples using reference compounds are shown below. However, it is noted that the carriers, surfactants and other additives which may be used in formulation of the insecticides
45 according to this invention are not limited by the following Formulation Examples. In the following Formulation Examples, "part" stands for "part by weight".

Formulation Example 1

50 32.5 Parts of the Compound No. 1 in Table 1, 3 parts of lignin sulfonic acid, 4 parts of polyoxyethylene alkylphenyl ether, 2 parts of silicon dioxide hydrate and 58.5 parts of clay are well mixed while being powdered to obtain a wettable powder.

Formulation Example 2

55

5.4 parts of the compound No. 2 in Table 1, 2 parts of silicon dioxide hydrate and 92.6 parts of talc are well mixed while being powdered to obtain a dust.

Formulation Example 3

5.4 parts of the compound No. 3 in Table 1, 3 parts of lignin sulfonate, 1 part of sodium dodecylbenzenesulfonate, 30 parts of bentonite and 60.6 parts of clay are well mixed while being powdered. The mixture is then kneaded with water, granulated and dried to obtain granules.

The insecticidal effects of the compounds of this invention are shown by referring to some Test Examples.


Test Example 1

Five rice seedlings having a height of about 7 cm were dipped, for 10 seconds, into aqueous dilutions of wettable powder each containing a predetermined concentration of each sample compound, prepared in accordance with the procedure as described in Formulation Example 1. After air drying, the roots of the five rice seedlings were wrapped with sanitary cotton dampened with water, and placed in a glass cylinder having a diameter of 3 cm and a height of 20 cm. Ten second instar larvae of a green rice leafhopper which had acquired resistance to chemicals were released into each glass cylinder. The cylinder was allowed to stand in a room maintained at 26°C. The mortality of the larvae was examined 48 hours after treatment. The results are shown in Table 2.

Table 2

Compound No.	Mortality (%), 500 ppm
5	100
6	100
7	100
8	100
9	100
10	100
11	100
Comparative Compound	70

Note: The comparative compound set forth in Table 2 is Sumithion (Trade Name), represented by the following formula.



Test Example 2

Five rice seedlings having a height of about 7 cm were dipped, for 10 seconds, into aqueous dilutions of wettable powder each containing a predetermined concentration of each sample compound, prepared in accordance with the procedure as described in Formulation Example 1. After air drying, the roots of the five rice seedlings were wrapped with sanitary cotton dampened with water, and placed in a glass cylinder having a diameter of 3 cm and a height of 20 cm. Ten second instar larvae of a brown rice planthopper which had acquired resistance to chemicals were released into each glass cylinder. The cylinder was allowed to stand in a room maintained at 26 °C. The mortality of the larvae was examined 48 hours after treatment. The results are shown in Table 3.

Table 3


Compound No.	Mortality (%), 500 ppm
5	100
6	100
7	100
8	100
9	100
10	100
11	100
Comparative Compound	65

Note: The comparative compound set forth in Table 3 is the same as that in Table 2.


As will be apparent from the foregoing, the compounds of this invention are extremely active for controlling various harmful insects and yet are low in toxicity to warm-blooded animals, fishes and Crustacea, the remaining quantity thereof after their use is small, and they do not produce phytotoxicity in various plants.

Claims

1. A hexahydrotriazine compound represented by the following general formula (I):




wherein R stands for C₁-C₈ alkyl or alkenyl of up to 8 carbon atoms; R₁ stands for C₁-C₈ alkyl, alkenyl of up to 8 carbon atoms or alkynyl of up to 8 carbon atoms; and R₂ stands for a group indicated by



2. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-pyridylmethyl)-3,5-dimethyl-2-nitroiminohexahydro-1,3,5-triazine.
3. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-thiazolylmethyl)-3,5-dimethyl-2-nitroiminohexahydro-1,3,5-triazine.
4. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-pyridylmethyl)-3-ethyl-5-methyl-2-nitroiminohexahydro-1,3,5-triazine.
5. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-pyridylmethyl)-3-propyl-5-methyl-2-nitroiminohexahydro-1,3,5-triazine.
6. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-thiazolylmethyl)-3-propyl-5-methyl-2-nitroiminohexahydro-1,3,5-triazine.
7. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-pyridylmethyl)-3-allyl-5-methyl-2-nitroiminohexahydro-1,3,5-triazine.
8. The hexahydrotriazine compound according to claim 1 which is 1-(2-chloro-5-pyridylmethyl)-3-propargyl-5-methyl-2-nitroiminohexahydro-1,3,5-triazine.
9. A process for preparing a hexahydrotriazine compound represented by the general formula (I) according to claim 1 comprising the step of reacting a compound represented by the general formula (II):



wherein R₁ and R₂ have the same meanings as defined in claim 1, with a compound represented by the general formula (III)



wherein R has the same meaning as defined in claim 1 and X stands for halogen.

10. A process for preparing a hexahydrotriazine compound represented by the general formula (I) according to claim 1 comprising the step of reacting a compound represented by the general formula (II):




wherein R₁ and R₂ have the same meanings as defined in claim 1, with a primary amine represented by the general formula (IV):




wherein R has the same meaning as defined in claim 1, and formalin or paraformaldehyde.


11. A process for preparing a hexahydrotriazine compound represented by the general formula (I) according to claim 1 comprising the step of reacting a compound represented by the general formula (V):



wherein R and R₁ have the same meanings as defined in claim 1 with a compound represented by the general formula (VI):




wherein R₂ has the same meaning as defined in claim 1 and Y stands for halogen or a group indicated by -OSO₃CH₃ or



12. An insecticide containing a suitable carrier, and, as an active ingredient, a hexahydrotriazine compound represented by the general formula (I) according to any one of claims 1 to 8.


40 **Patentansprüche**

1. Hexahydrotriazinverbindung der nachstehenden allgemeinen Formel (I):




in der R einen C₁-C₈-Alkylrest oder Alkenylrest mit bis zu 8 Kohlenstoffatomen bedeutet; R₁ einen C₁-C₈-Alkylrest, Alkenylrest mit bis zu 8 Kohlenstoffatomen oder Alkinyrest mit bis zu 8 Kohlenstoffatomen

bedeutet, und R₂ eine durch




dargestellte Gruppe bedeutet.

- 10 2. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-pyridylmethyl)-3,5-dimethyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
- 15 3. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-thiazolylmethyl)-3,5-dimethyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
4. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-pyridylmethyl)-3-ethyl-5-methyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
- 20 5. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-pyridylmethyl)-3-propyl-5-methyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
6. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-thiazolylmethyl)-3-propyl-5-methyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
- 25 7. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-pyridylmethyl)-3-allyl-5-methyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
8. Hexahydrotriazinverbindung gemäß Anspruch 1, die 1-(2-Chlor-5-pyridylmethyl)-3-propargyl-5-methyl-2-nitroimino-
nohexahydro-1,3,5-triazin ist.
- 30 9. Verfahren zur Herstellung einer Hexahydrotriazinverbindung der allgemeinen Formel (I) gemäß Anspruch 1, umfassend den Schritt der Umsetzung einer Verbindung der allgemeinen Formel (II):




in der R₁ und R₂ dieselben Bedeutungen wie in Anspruch 1 haben, mit einer Verbindung der allgemeinen Formel (III):




in der R dieselbe Bedeutung wie in Anspruch 1 hat, und X ein Halogenatom bedeutet.

- 50 10. Verfahren zur Herstellung einer Hexahydrotriazinverbindung der allgemeinen Formel (I) gemäß Anspruch 1, umfassend den Schritt der Umsetzung einer Verbindung der allgemeinen Formel (II):




in der R_1 und R_2 dieselben Bedeutungen wie in Anspruch 1 haben, mit einem primären Amin der allgemeinen Formel (IV):




in der R dieselbe Bedeutung wie in Anspruch 1 hat, und Formalin oder Paraformaldehyd.


11. Verfahren zur Herstellung einer Hexahydrotriazinverbindung der allgemeinen Formel (I) gemäß Anspruch 1, umfassend den Schritt der Umsetzung einer Verbindung der allgemeinen Formel (V):



in der R und R_1 dieselben Bedeutungen wie in Anspruch 1 haben, mit einer Verbindung der allgemeinen Formel (VI):



in der R_2 dieselbe Bedeutung wie in Anspruch 1 hat, und Y ein Halogenatom oder eine durch $-\text{OSO}_2\text{CH}_3$ oder




dargestellte Gruppe bedeutet.


12. Insektizid, enthaltend einen geeigneten Träger und als Wirkstoff eine Hexahydrotriazinverbindung der allgemeinen Formel (I) gemäß einem der Ansprüche 1 bis 8.

Revendications

1. Dérivé d'hexahydrotriazine représenté par la formule générale (I) suivante :




20 dans laquelle R représente un groupe alkyle en C₁-C₈ ou un groupe alcényle ayant jusqu'à 8 atomes de carbone ; R₁ représente un groupe alkyle en C₁-C₈, un groupe alcényle ayant jusqu'à 8 atomes de carbone ou un groupe alcyne ayant jusqu'à 8 atomes de carbone ; et R₂ représente un groupe représenté par



- 30 2. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-pyridylméthyl)-3,5-diméthyl-2-nitro-iminohexahydro-1,3,5-triazine.
3. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-thiazolylméthyl)-3,5-diméthyl-2-nitro-iminohexahydro-1,3,5-triazine.
- 35 4. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-pyridylméthyl)-3-éthyl-5-méthyl-2-nitro-iminohexahydro-1,3,5-triazine.
5. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-pyridylméthyl)-3-propyl-5-méthyl-2-nitro-iminohexahydro-1,3,5-triazine.
- 40 6. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-thiazolylméthyl)-3-propyl-5-méthyl-2-nitroiminohexahydro-1,3,5-triazine.
7. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-pyridylméthyl)-3-allyl-5-méthyl-2-nitro-iminohexahydro-1,3,5-triazine.
- 45 8. Dérivé d'hexahydrotriazine selon la revendication 1, qui est la 1-(2-chloro-5-pyridylméthyl)-3-propargyl-5-méthyl-2-nitroiminohexahydro-1,3,5-triazine.
- 50 9. Procédé pour préparer un dérivé d'hexahydrotriazine représenté par la formule générale (I) selon la revendication 1 comprenant l'étape de réaction d'un composé représenté par la formule générale (II) :



dans laquelle R_1 et R_2 sont tels que définis dans la revendication 1, avec un composé représenté par la formule générale (III)



dans laquelle R est tel que défini dans la revendication 1 et X représente un atome d'halogène.

10. Procédé de préparation d'un dérivé d'hexahydrotriazine représenté par la formule générale (I) selon la revendication 1 comprenant l'étape de réaction d'un composé représenté par la formule générale (II) :




dans laquelle R_1 et R_2 sont tels que définis dans la revendication 1, avec une amine primaire représentée par la formule générale (IV) :




dans laquelle R est tel que défini dans la revendication 1, et la formaline ou le paraformaldéhyde.


11. Procédé de préparation d'un dérivé d'hexahydrotriazine représenté par la formule générale (I) selon la revendication 1 comprenant l'étape de réaction d'un composé représenté par la formule générale (V) :



dans laquelle R et R_1 sont tels que définis dans la revendication 1, avec un composé représenté par la formule générale (VI) :



dans laquelle R_2 est tel que défini dans la revendication 1 et Y représente un atome d'halogène ou un groupe de formule $-\text{OSO}_2\text{CH}_3$ ou



5

12. Insecticide contenant un support approprié et, en tant qu'ingrédient actif, un dérivé d'hexahydrotriazine représenté par la formule générale (I) selon l'une quelconque des revendications 1 à 8.

10

15

20

25

30

35

40

45

50

55